


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C07D 487/04, A61K 31/50, 31/41, A61P 29/00 // (C07D 487/04, 233:00, 241:00)</b>		<b>A1</b>	(11) International Publication Number: <b>WO 00/35921</b>
		(43) International Publication Date: 22 June 2000 (22.06.00)	
(21) International Application Number: PCT/EP99/09806 (22) International Filing Date: 11 December 1999 (11.12.99) (30) Priority Data: 60/112,653 17 December 1998 (17.12.98) US (71) Applicant: F. HOFFMANN-LA ROCHE AG [CH/CH]; 124 Grenzacherstrasse, CH-4070 Basle (CH). (72) Inventors: LUK, Kin-Chun; 66 Evergreen Drive, North Caldwell, NJ 07006-4622 (US). MICHOD, Christophe; Apartment 2A, 411 East 87th Street, New York, NY 10128 (US). (74) Agent: LOESCHNER, Thomas; 124 Grenzacherstrasse, CH-4070 Basle (CH).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: 4,5-PYRAZINOXINDOLES AS PROTEIN KINASE INHIBITORS			
<div style="text-align: center;">  <p>(I)</p> </div>			
(57) Abstract 4,5-pyrazinoxindoles having formula (I), inhibit or modulate protein kinases, in particular JNK protein kinases and are useful as anti-inflammatory agents, particularly in the treatment of rheumatoid arthritis.			

What Is Claimed Is:

1. A compound of formula

5



wherein:

$R^1$  and  $R^2$  are independently selected from the group consisting of

- hydrogen,  
 10 -OR<sup>4</sup>,  
 -COR<sup>4</sup>,  
 -COOR<sup>4</sup>,  
 -CONR<sup>5</sup>R<sup>6</sup>,  
 -NR<sup>5</sup>R<sup>6</sup>,

15 lower alkyl which may be substituted by a member of the group (a) consisting of -OR<sup>4</sup>, -NR<sup>5</sup>R<sup>6</sup>, halogen, -COR<sup>4</sup>, -COOR<sup>4</sup>, -OCOR<sup>4</sup>, -CONR<sup>5</sup>R<sup>6</sup>, -CN, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>; or by cycloalkyl, heterocycle, aryl, and heteroaryl, wherein the cycloalkyl and heterocycle each may be substituted by the group R<sup>11</sup> and the aryl and heteroaryl each may be substituted by the group R<sup>12</sup>;

20 cycloalkyl which may be substituted by a member of the group (a) a defined earlier, or by lower alkyl, heterocycle, aryl, and heteroaryl, wherein the lower alkyl and heterocycle each may be substituted by the group R<sup>11</sup> and the aryl and heteroaryl each may be substituted by the group R<sup>12</sup>;

heterocycle which may be substituted by a member of the group (a) as  
 25 defined earlier, or by lower alkyl, cycloalkyl, aryl, and heteroaryl, wherein the lower alkyl and cycloalkyl each may be substituted by the group R<sup>11</sup> and the aryl and heteroaryl each may be optionally substituted by the group R<sup>12</sup>;

aryl which may be substituted by a member of the group (b) consisting of -OR<sup>4</sup>, -NR<sup>5</sup>R<sup>6</sup>, halogen, -NO<sub>2</sub>, perfluoroalkyl, -COR<sup>4</sup>, -COOR<sup>4</sup>, -OCOR<sup>4</sup>, -CONR<sup>5</sup>R<sup>6</sup>, -CN, -  
 30 SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>; or by lower alkyl, cycloalkyl, heterocycle, aryl, and heteroaryl, and wherein the lower alkyl, cycloalkyl and heterocycle each may be substituted by the group R<sup>11</sup> and the aryl and heteroaryl each may be substituted by the group R<sup>12</sup>,

- heteroaryl which may be substituted by a member of the group (b) as defined earlier, or by lower alkyl, cycloalkyl, heterocycle, aryl, and heteroaryl and wherein the lower alkyl, cycloalkyl and heterocycle each may be optionally substituted by the group  $R^{11}$  and the aryl and heteroaryl each may be substituted by the group  $R^{12}$ , or
- 5 alternatively,  $R^1$  and  $R^2$  can form a ring having 5-7 atoms, said ring optionally including one or more heteroatoms and being optionally substituted by a member of the group consisting of  $-OR^8$ ,  $-COR^7$ ,  $-COOR^7$ ,  $-OCOR^4$ ,  $-CONR^7R^9$ ,  $-NR^8R^9$ , or lower alkyl which may be substituted by the group  $R^{11}$ ;
- 10  $R^3$  is hydrogen,  $-OR^4$ ,  $-COR^4$ ,  $-COOR^4$ ,  $-OCOR^4$ ,  $-CONR^5R^6$ , halogen,  $-CN$ , perfluoroalkyl  $-NR^5R^6$ , or lower alkyl which may be substituted by  $-OR^4$ ,  $-OCOR^4$ , or  $-NR^5R^6$ ;
- $R^4$  is hydrogen,
- 15 lower alkyl which may be substituted by a member of the group (c) consisting of  $-OR^8$ ,  $-COOR^7$ ,  $-COR^7$ ,  $-CONR^5R^6$ ,  $-NR^5R^6$ ,  $-SO_2R^7$ ,  $-SO_2NR^5R^6$ ; or by cycloalkyl, heterocycle, aryl, and heteroaryl, and wherein the cycloalkyl and heterocycle each may be substituted by the group  $R^{11}$  and the aryl and heteroaryl each may be substituted by the group  $R^{12}$ ,
- 20 cycloalkyl which may be substituted by a member of the group (c) or by lower alkyl, heterocycle, aryl, and heteroaryl, and wherein the lower alkyl and heterocycle each may be substituted by the group  $R^{11}$  and the aryl and heteroaryl each may be substituted by the group  $R^{12}$ ,
- heterocycle which may be substituted by a member of the group (c) or by
- 25 cycloalkyl, lower alkyl, aryl, and heteroaryl, and wherein the cycloalkyl and lower alkyl each may be substituted by the group  $R^{11}$  and the aryl and heteroaryl each may be substituted by the group  $R^{12}$ ,
- aryl which may be substituted by a member of the group (d) consisting of  $-OR^8$ ,  $-COOR^7$ ,  $-COR^7$ ,  $-CONR^5R^6$ ,  $-NR^5R^6$ ,  $-NO_2$ , halogen, perfluoroalkyl,  $-SO_2R^7$ ,  $-SO_2NR^5R^6$ ; or by lower alkyl, cycloalkyl, heterocycle, aryl, and heteroaryl, and wherein the
- 30 lower alkyl, cycloalkyl and heterocycle each may be substituted by the group  $R^{11}$  and the aryl and heteroaryl each may be substituted by the group  $R^{12}$ , and
- heteroaryl which may be substituted by a member of the group (d) or by
- 35 cycloalkyl, lower alkyl, heterocycle, aryl, and heteroaryl, and wherein the lower alkyl,

cycloalkyl and heterocycle each may be substituted by the group  $R^{11}$  and the aryl and heteroaryl each may be substituted by the group  $R^{12}$ ;

$R^5$  and  $R^6$  are each independently

- 5                   hydrogen,  
                  -COR<sup>7</sup>,  
                  -COOR<sup>7</sup>,  
                  -CONR<sup>7</sup>R<sup>9</sup>,

- lower alkyl which may be substituted by a member of the group (e)  
10   consisting of -OR<sup>8</sup>, -COOR<sup>7</sup>, -COR<sup>7</sup>, -CONR<sup>7</sup>R<sup>8</sup>, -NR<sup>7</sup>R<sup>8</sup>, -SO<sub>2</sub>R<sup>7</sup>, -SO<sub>2</sub>NR<sup>7</sup>R<sup>8</sup>; or by  
                  cycloalkyl, heterocycle, aryl, and heteroaryl, and wherein the cycloalkyl and heterocycle  
                  each may be substituted by the group  $R^{11}$  and the aryl and heteroaryl each may be  
                  substituted by the group  $R^{12}$ ,

- cycloalkyl which may be substituted by a member of the group (e) as  
15   defined earlier, or by lower alkyl, heterocycle, aryl, and heteroaryl, and wherein the lower  
                  alkyl and heterocycle each may be substituted by the group  $R^{11}$  and the aryl and heteroaryl  
                  each may be substituted by the group  $R^{12}$ ,

- heterocycle which may be substituted by a member of the group (e) as  
                  defined earlier, or by cycloalkyl, lower alkyl, aryl, and heteroaryl, and wherein the  
20   cycloalkyl and lower alkyl each may be substituted by the group  $R^{11}$  and the aryl and  
                  heteroaryl each may be substituted by the group  $R^{12}$ ,

- aryl which may be substituted by a member of the group (f) consisting of  
                  OR<sup>8</sup>, -COOR<sup>7</sup>, -COR<sup>7</sup>, -CONR<sup>7</sup>R<sup>8</sup>, -NR<sup>7</sup>R<sup>8</sup>, -NO<sub>2</sub>, halogen, perfluoroalkyl, -SO<sub>2</sub>R<sup>7</sup>,  
                  -SO<sub>2</sub>NR<sup>7</sup>R<sup>8</sup>; or by lower alkyl, cycloalkyl, heterocycle, aryl, and heteroaryl, and wherein the  
25   lower alkyl, cycloalkyl and heterocycle each may be substituted by the group  $R^{11}$  and the  
                  aryl and heteroaryl each may be substituted by the group  $R^{12}$ , and

- heteroaryl which may be substituted by a member of the group (f) as  
                  defined earlier, or by lower alkyl, cycloalkyl, heterocycle, aryl, and heteroaryl, and wherein  
                  the lower alkyl, cycloalkyl and heterocycle each may be substituted by the group  $R^{11}$  and the  
30   aryl and heteroaryl each may be substituted by the group  $R^{12}$ ; or alternatively,  
                  -NR<sup>5</sup>R<sup>6</sup> can form a ring having 3 to 7 atoms, said ring optionally including one or more  
                  additional hetero atoms and being optionally substituted by lower alkyl, -OR<sup>8</sup>,  
                  -COR<sup>7</sup>, -COOR<sup>7</sup>, -CONR<sup>7</sup>R<sup>9</sup>, or -NR<sup>8</sup>R<sup>9</sup>;

- 35                    $R^7$  is hydrogen or lower alkyl which may be substituted by a member of the group  
                  consisting of cycloalkyl, heterocycle, aryl, heteroaryl, -OR<sup>9</sup>, or -NR<sup>8</sup>R<sup>9</sup>;

$R^8$  is hydrogen,  $-\text{COR}^9$ ,  $-\text{CONR}^{10}\text{R}^9$ , or lower alkyl which may be substituted by  $R^{11}$ ;

$R^9$  and  $R^{10}$  are each independently hydrogen or lower alkyl;

5  $R^{11}$  is  $-\text{OR}^9$ ,  $-\text{COR}^9$ ,  $-\text{COOR}^9$ ,  $-\text{OCOR}^9$ ,  $-\text{CONR}^9\text{R}^{10}$ ,  $-\text{NR}^9\text{R}^{10}$ ,  $-\text{N}(\text{COR}^9)\text{R}^{10}$ ,  $-\text{SO}_2\text{R}^9$ , or  $-\text{SO}_2\text{NR}^9\text{R}^{10}$ ;

10  $R^{12}$  is  $-\text{OR}^9$ ,  $-\text{COR}^9$ ,  $-\text{COOR}^9$ ,  $-\text{OCOR}^9$ ,  $-\text{CONR}^9\text{R}^{10}$ ,  $-\text{NR}^9\text{R}^{10}$ ,  $-\text{N}(\text{COR}^9)\text{R}^{10}$ ,  $-\text{SO}_2\text{R}^9$ ,  $-\text{SO}_2\text{NR}^9\text{R}^{10}$ , halogen,  $-\text{CN}$ ,  $-\text{NO}_2$ , or perfluoroalkyl; and

X is -N- or -C-.

and prodrugs and pharmaceutically active metabolites of compounds of Formula I; and the pharmaceutically acceptable salts of the foregoing compounds.

15

2. A compound of claim 1, wherein

$R^1$  and  $R^2$  are independently

hydrogen,

$-\text{NR}^5\text{R}^6$ ,

20

lower alkyl which may be substituted by  $R^{11}$ , cycloalkyl, heterocycle, aryl and heteroaryl, wherein the cycloalkyl and heterocycle may be substituted by  $R^{11}$ , and the aryl and heteroaryl may be substituted by  $R^{12}$ ;

cycloalkyl which may be substituted by  $R^{11}$ , lower alkyl, heterocycle, aryl and heteroaryl, wherein the lower alkyl and heterocycle may be substituted by  $R^{11}$ , and  
25 the aryl and heteroaryl may be substituted by  $R^{12}$ ;

heterocycle which may be substituted by  $R^{11}$ , lower alkyl, cycloalkyl, aryl and heteroaryl, wherein the lower alkyl and cycloalkyl may be substituted by  $R^{11}$ , and the aryl and heteroaryl may be substituted by  $R^{12}$ ;

30 aryl which may be substituted by  $R^{12}$ , lower alkyl, cycloalkyl, heterocycle, aryl, and heteroaryl, wherein the lower alkyl, heterocycle and cycloalkyl may be substituted by  $R^{11}$ , and the aryl and heteroaryl may be substituted by  $R^{12}$ ;

heteroaryl which may be substituted by  $R^{12}$ , lower alkyl, cycloalkyl, heterocycle, aryl, and heteroaryl, wherein the lower alkyl, cycloalkyl and heterocycle may be substituted by  $R^{11}$ , and the aryl and heteroaryl may be substituted by  $R^{12}$ ; or alternatively,

R<sup>1</sup> and R<sup>2</sup> may form a ring having 5 to 7 atoms and optionally being substituted by the group consisting of -OR<sup>8</sup>, -COR<sup>7</sup>, -COOR<sup>7</sup>, -CONR<sup>7</sup>R<sup>9</sup>, -NR<sup>8</sup>R<sup>9</sup>, and lower alkyl which may be substituted by R<sup>11</sup>.

- 5     3.     The compound of claim 2 wherein R<sup>3</sup> is hydrogen, -OR<sup>4</sup>, -NR<sup>5</sup>R<sup>6</sup>, or lower alkyl which may be substituted by the group consisting of -OR<sup>4</sup> and -NR<sup>5</sup>R<sup>6</sup>.
4.     The compound of claim 2 wherein R<sup>3</sup> is hydrogen, -OR<sup>9</sup>, or lower alkyl which may be substituted by the group consisting of -OR<sup>9</sup> and -NR<sup>9</sup>R<sup>10</sup>.
- 10     5.     The compound of claim 1, which is (Z)-7,9-Dihydro-2,3-dimethyl-9-[(3-methoxy-1H-pyrrol-2-yl)methylene]-8H-pyrrolo-[3,2-f]quinoxalin-8-one
6.     The compound of claim 1, which is (Z)-3-Butyl-7,9-dihydro-9-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-methyl-8H-pyrrolo[3,2-f]quinoxalin-8-one
- 15     7.     The compound of claim 1, which is (Z)-2-butyl-7,9-dihydro-9-[(3-methoxy-1H-pyrrol-2-yl)methylene]-3-methyl-8H-pyrrolo[3,2-f]quinoxalin-8-one
- 20     8.     The compound of claim 1, which is (Z)-7,9-Dihydro-9-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2-methyl-3-phenyl-8H-pyrrolo[3,2-f]quinoxalin-8-one
9.     The compound of claim 1, which is (Z)-7,9-dihydro-9-[(3-methoxy-1H-pyrrol-2-yl)methylene]-3-methyl-2-phenyl-8H-pyrrolo[3,2-f]quinoxalin-8-one
- 25     10.    The compound of claim 1, which is (Z)-7,9-Dihydro-2,3-di-(2-furanyl)-9-[(3-methoxy-1H-pyrrol-2-yl)methylene]-8H-pyrrolo[3,2-f]quinoxalin-8-one
11.    The compound of claim 1, which is (Z)-1,3,5,6,7,8-Hexahydro-3-[(3-methoxy-1H-pyrrol-2-yl)methylene]-2H-pyrrolo[3,2-a]phenazin-2-one
- 30     12.    A pharmaceutical composition comprising as an active ingredient a compound of any one of claims 1 to 11 and a pharmaceutically acceptable carrier or excipient.

13. A compound of any one of claims 1 to 11 for use as a medicament, particularly for the treatment and/or control of inflammation and neurodegenerative diseases, particularly rheumatoid arthritis, or for treating solid tumors, in particular breast or colon tumors.

5 14. The use of a compound of formula I or a pharmaceutically acceptable salt thereof as defined in any one of claims 1 to 11 in the preparation of a medicament containing such compound for the treatment and/or control of inflammation and neurodegenerative diseases, particularly rheumatoid arthritis, or for treating solid tumors, in particular breast or colon tumors

10

15. The invention as described hereinbefore.

International Application No  
PCT/EP 99/09806

1



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/09806

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>WO 96 32380 A (PHARMACIA S.P.A.)  17 October 1996 (1996-10-17)  cited in the application  claims 1-10</p>	1-15

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No  
PCT/EP 99/09806

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9915500 A	01-04-1999	AU 9740798 A	12-04-1999
WO 9910325 A	04-03-1999	AU 9158498 A	16-03-1999
WO 9725986 A	24-07-1997	AU 708167 B	29-07-1999
		AU 1398697 A	11-08-1997
		AU 702045 B	11-02-1999
		AU 1398797 A	11-08-1997
		CA 2214744 A	24-07-1997
		CA 2214759 A	24-07-1997
		EP 0815859 A	07-01-1998
		EP 0816338 A	07-01-1998
		EP 9800757 A	28-07-1998
		HU 9726242 A	24-07-1997
		WO 974280 A	11-11-1997
		NO 5977130 A	02-11-1999
		US 5965600 A	12-10-1999
WO 9807695 A	26-02-1998	AU 4155697 A	06-03-1998
		EP 0929520 A	21-07-1999
WO 9632380 A	17-10-1996	EP 0764152 A	26-03-1997
		JP 10501821 T	17-02-1998
		US 5849710 A	15-12-1998